

Similarities of Akt activation in cutaneous wound repair and skin carcinogenesis

Nesrine I. Affara¹, Brandon L. Schanbacher², Ping Pei³, Susan R. Mallery³, Carol S. Trempus⁴, John A. Bauer², and Fredika M. Robertson¹

¹Department of Molecular Virology, Immunology & Medical Genetics, The Ohio State University, Columbus, Ohio;

²The Center for Cardiovascular Pharmacology, Columbus Children's Research Institute, Columbus, Ohio;

³Department of Oral Maxillofacial Surgery, Oral Pathology and Periodontology, The Ohio State University, Columbus, Ohio; ⁴National Center for Toxicogenomics, National Institute of Environmental Health Sciences, Research Triangle Park, NC

Nrf transcription factors are crucial for skin tumor prevention but not for wound healing

Ulrich auf dem Keller¹, Marcel Huber², Tobias Beyer¹, Philippe Bugnon¹, Susanne Braun¹, Thomas Rüllicke³, Delinda A. Johnson⁴, Jeffrey A. Johnson⁴, Daniel Hohl², and Sabine Werner¹

¹Institute of Cell Biology, Department of Biology, ETH Zürich, Hönggerberg, Zürich, Switzerland; ²Service de Dermatologie, CHUV Hôpital Beaumont, Lausanne, Switzerland; ³Institute of Laboratory Animal Science, Universität Zürich, Zürich, Switzerland; ⁴University of Wisconsin, School of Pharmacy, Madison, WI

Bone marrow-derived cells fuse with normal and transformed intestinal progenitor cells

Adnan Z. Rizvi¹, John R. Swain², Paige S. Davies², Alexis Bailey³, Adria D. Decker², Holger Willenbring⁴⁺, Markus Grompe⁴⁺, William H. Fleming³⁺, and Melissa H. Wong²⁺

¹ Department of Surgery, ²Departments of Dermatology and of Cell and Developmental Biology, ⁴Department of Molecular and Medical Genetics, ³Center for Hematologic Malignancies, Division of Hematology and Medical Oncology, +Oregon Cancer Institute, Oregon Stem Cell Center, Oregon Health & Science University, Portland, OR

Evaluations of a novel Topical Plant Extract (TPE)* therapy on second-degree burns and Methicillin-Resistant Staphylococcus aureus (MRSA) using porcine wound models

Stephen C. Davis, Ramon Montero, Alex Cazzaniga and Patricia Mertz

University of Miami Department of Dermatology & Cutaneous Surgery, Miami, FL

Regulation of VEGF-A protein activity during wound repair

Detlev Roth¹, Michael Piekarek¹, Mats Paulsson^{2,3}, Thomas Krieg^{1,3} Jeffrey M. Davidson^{4,5}, Sabine A. Eming¹

¹Department of Dermatology, University of Cologne, Cologne, Germany; ²Center for Biochemistry, Medical Faculty, University of Cologne, Cologne, Germany; ³Center for Molecular Medicine, University of Cologne, Cologne, Germany; ⁴Department of Pathology, Vanderbilt University School of Medicine and ⁵VA Medical Center, Nashville, TN

Effect of elevated polyamine biosynthesis in cutaneous wound healing

Candace Sewter¹, Karen DeFeo¹, Rebecca Morris², and Susan Gilmour¹

¹Lankenau Institute for Medical Research, Wynnewood, PA, ²Columbia University, New York, NY

Fibroblast Growth Factor Receptor 2-IIIb (Fgfr2b) is a critical mediator of intercellular signalling in development and disease

Richard Grose

Department of Tumour Biology, Cancer Research UK Clinical Centre, Bart's & The London School of Medicine & Dentistry, London, UK

Why won't this wound heal??

Carolyn Hale^{1,2}, Kevin White¹

¹Department of Dermatology, Oregon Health and Science University, Portland, OR; ² Private Practice, Keizer, OR

Overexpression of smad7 in keratinocytes accelerates cutaneous wound healing

Gangwen Han, Allen. G. Li, Philip Owens, Xiao-Jing Wang

Depts of Otolaryngology, Dermatology, Cell & Developmental Biology, Oregon Health & Science University, Portland, OR

p51/p63 promotes lactadherin (MFG-E8) expression in epithelia.

Takeshi Okuyama, Iyoko Katoh, Yoshiya Tomimori, Shingo Sato, Shun-ichi Kurata, Shizuko Kobayashi, and Yoji Ikawa

Ikawa Laboratory, RIKEN, Wako, Japan

Reflection-mode confocal microscopy sensitive to ultrastructure of skin and other biological tissues

Steven L. Jacques¹, Daniel S. Gareau¹, James Lagowski¹, Glenn Merlino², Molly Kulesz-Martin¹

¹Oregon Health & Science University, Portland, OR; ²National Cancer Institute, NIH

p63 isoform switching is required for induction of IKK α and epidermal morphogenesis

Maranke I. Koster¹, Antonio Costanzo³, Yuji Sano⁴, Michael Karin⁴ and Dennis R. Roop^{1,2}

¹Departments of Molecular and Cellular Biology and ²Dermatology, Baylor College of Medicine, Houston, TX; ³Department of Dermatology, University of Rome "Tor Vergata," Rome, Italy; ⁴Department of Pharmacology, University of California, San Diego, CA

Analysis of caspase expression and activity in healing wounds

Melanie K. Kuechle, Tamara Brockow, Robert Underwod, Marcia Usui, John Olerud

University of Washington School of Medicine Department of Medicine, Division of Dermatology, Seattle, WA

EGF Modulates Slug Expression During Cutaneous Wound Reepithelialization

Donna F. Kusewitt, Changsun Choi, Laurie G. Hudson

The Ohio State University College of Veterinary Medicine, Columbus, OH (DFK, CC), The University of New Mexico College of Pharmacy, Albuquerque, NM (LGH)

Fibroblast senescence, cytokine production, microbial colonization and chronic wounds

Jonathan Mansbridge, Chris Loryman

Smith and Nephew, La Jolla, CA

TGF β 1 overexpression in mouse keratinocytes delays cutaneous wound healing

Philip Owens, Allen G. Li, Gangwen Han, Xiao-Jing Wang

Depts of Otolaryngology, Dermatology, Cell & Developmental Biology, Oregon Health & Science University, Portland, OR

Genetic analyses reveal potential roles for novel IL1 and IL1R family members in skin inflammation

Jacques Peschon¹, Huyen Dinh¹, David Kugler¹, Esther Trueblood² and Hal Blumberg¹

Departments of ¹Inflammation and ²Pathology, Amgen Washington, Seattle, WA

Can a topical antimicrobial agent penetrate a bi-layered cell therapy and be effective against Methicillin-Resistant Staphylococcus aureus (MRSA)?

Franco Pissani, Navid Bouzari, Ramon B Montero, Stephen C Davis

University of Miami School of Medicine, Department of Dermatology & Cutaneous Surgery, Miami, FL

The role of melatonin on scarring in an incisional model of dermal wound healing in rats

K Pugazhenti, M Kapoor, P Young, A Clarkson, I Hall, I Appleton

Department of Pharmacology and Toxicology, University of Otago, Dunedin, New Zealand

β 2-adrenergic receptor antagonists are pro-motogenic in keratinocytes: Potential therapy for promoting wound re-epithelialization

Christine E. Pullar and R. Rivkah Isseroff

Department of Dermatology, University of California, Davis School of Medicine, Davis, CA

The role of RIP4 in squamous epithelial development

Ryan B. Rountree, Cynthia R. Willis, Jacques Peschon, and Pam Holland

Amgen Washington, AW2-D2242, 1201 Amgen Court W., Seattle, WA 98119

Elevated expression of both β 1 and β 4 integrins in slow cycling mouse corneal epithelial cells

A. Pajooheh-Ganji^{a,b,c}, S. Pal-Ghosh^{a,c}, S.J. Simmens^{c,e}, and M.A. Stepp^{a,d,e,*}

^aDepartment of Anatomy and Cell Biology, ^bDepartment of Biological Sciences,

^cDepartment of Epidemiology and Biostatistics, ^dDepartment of Ophthalmology

^eThe George Washington University Medical Center, Washington, DC

Protein expression in the epidermis of non-healing ulcers from patients with diabetes

Usui ML, Carter WG*†, Underwood RA, Olerud JE

Departments of Medicine (Dermatology) and Pathobiology*, University of Washington, Seattle, WA and †Fred Hutchinson Cancer Research Center, Seattle, WA

Non-endothelial effects of vascular endothelial growth factor: Implications for wound repair

Traci A. Wilgus and Luisa A. DiPietro

Loyola University Medical Center, Burn & Shock Trauma Institute, Department of Surgery, Maywood, IL